

Exploring the Disease-causing Potential of *Aeromonas Hydrophila* Obtained from Freshwater Carp - *Cirrhinus Mrigala*

Ravi D. Barde

Department of Zoology, SGB College, Purna, Maharashtra, India,
ravibarde4u@gmail.com

Abstract

In India, fish bacteriology has a slow progress when compared to developed countries. The bacterial diseases reported so far are dropsy or hemorrhagic septicemia, fin rot, ulcer disease, columnaris disease, eye disease of Catla, skin disease of Catfish, streptococcal infection etc. Among the Indian Major Carps are *Labeo calbasu*, *Cirrhinus mrigala*, *Cyprinus carpio*, *Catla catla* and *Labeo rohita*. *Cyprinus carpio* and *Cirrhinus mrigala* are the second fast growing and are widely consumed throughout India.

Cirrhinus mrigala is susceptible to bacterial diseases in natural as well as cultured system and is infected with *Aeromonas hydrophila*. The survey revealed the occurrence in the region of Marathwada in most of the waterbodies surveyed. Two most virulent isolate among the isolated strains were tested for their ability to cause symptoms and subsequent death in the experimental fishes. The preliminary tests indicated that ulcers were associated with *Aeromonas hydrophila* AH-04 and AH-07 isolates. The LD₅₀ values were estimated as 0.837×10^5 , and 0.878×10^5 for isolates AH-04 and AH-07, respectively

Keywords: *Cirrhinus mrigala*, *Aeromonas hydrophila*, LD 50.

INTRODUCTION

Diseases caused by bacterial are associated with substantial death in both wild and cultured fishes. The authentic part of these pathogenic microorganisms may vary from a prime pathogen to that of an opportunistic pathogen that imparts its host decline by initiating a disease process (Barde, 2021).

Infection in fishes brings about deleterious effects in their growth, resistance to stress and fecundity, and in addition, they are vulnerable to predation and sub optimal environmental factors. All these variables decimate the fish, indirectly impinging a heavy economic loss to the fish farmers, and it becomes imperative for them to control the infection by constant surveillance and development of adequate control measures to prevent spreading of diseases and to promote healthy environment in all the farms.

It is common knowledge that cultivable fishes are susceptible to disease-causing pathogenic bacteria (Darak & Barde, 2014; Darak et al., 2021) and fungi (Barde et al., 2020, 2021). Bacterial diseases of fish have always caused night mares among fish farmers. Among bacterial diseases, furunculosis, ulcer disease, columnaris disease and several others have caused epizootics of such magnitude as to threaten the entire salmonid fisheries of the affected regions. Furunculosis of salmonid fish received more attention from pathologists, and has been a serious problem in hatcheries of many countries, like USA, Germany, Italy, Canada and UK. It is the first bacterial fish disease which has been worked out fully.

In India, fish bacteriology has a slow progress when compared to developed countries. The bacterial diseases reported so far are dropsy or hemorrhagic septicemia, fin rot, ulcer disease,

Co-ordinator
IQAC

Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn) Dist. Parbhani - 431511 (M.S.)



1725
PRINCIPAL

Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn.) Dist. Parbhani

columnaris disease, eye disease of Catla, skin disease of Catfish, streptococcal infection etc (Darak and Barde, 2015). Therefore, it is necessary to throw emphasis on induced infections and to develop suitable control methods.

The Indian major carps are the most important food fishes and they are of very high commercial value. They are widely cultivated all over India, especially in central and South India and particularly in Tamil nadu. Private entrepreneurs have undertaken large scale fish farming by converting many a rice field into commercially viable fish farms. Among the Indian Major Carps are *Labeo calbasu*, *Cyprinus carpio*, *Cirrhinus mrigala*, *Catla catla* and *Labeo rohita*. *Cyprinus carpio* and *Cirrhinus mrigala* are the second fast growing and are widely consumed throughout India. *Cirrhinus mrigala* is susceptible to bacterial diseases in natural as well as cultured system. Hence, it was chosen as the experimental organism to find out the effects of pathogenic bacterial strain on the physiology and nutritional value of the fishes following infection by bacterial pathogen.

Material and methodology:

Fish Samples

Fish samples of fish species *Cirrhinus mrigala* species had been collected typically from surrounding natural bodies of water and cultivated systems. After determining the source, some fish were also bought from the local markets for this investigation. Samples were taken for analysis after processing at the laboratory immediately. Immediately after catching with sterilized cotton swabs, samples of different body parts were obtained. Within 4 hours of selection the samples were cultivated. The fishes were selected and considering their

easier availability, the health status of the fishes was given high priority.

Isolation and Maintenance of cultures

The isolation of heterotrophic, aerobic and anaerobic bacterial communities in terms of cfu/ml were identified and enumerated by the standard methods described by Cheesbrough (1989) and Bergey's Manual for Systematic Bacteriology (1986). Cultures were preserved by aseptically transferring the bacterial cultures to freshly prepared sterile nutrient agar slants after every three weeks. The stock cultures were stored at 4°C. For experimental works, subcultures were made from the stock cultures in suitable media before use. The cultures were also examined at regular intervals to test their pathogenicity.

Determination of LDW values of the pathogenic bacteria

The two pathogenic bacteria, *Aeromonas* spp. (AH04 and AH07) were grown in Brain Heart Infusion Broth at 30°C for 48 hours. These cultures were used to make five 10 fold dilutions of calculated by the spread plate method. 0.05 ml of each of these diluted cell suspensions were injected intraperitoneally to 10 healthy fishes. The doses received by the fishes ranged from 1×10^3 to 1×10^7 cfu. Each of the 10 control fishes received 0.05 ml of sterile 0.85% NaCl solution. Following injection, the fishes were observed for a 15 days period to record the appearance of disease symptoms and mortality. The dead fishes were immediately sacrificed and parts of liver and kidney were incubated in nutrient broth for re-isolation of the bacteria. The lethal dose 50% end, (LD50) was calculated from the relationship between the probits of percentage mortalities and the logs of the dilution series of bacterial suspension.


Co-ordinator
IQAC
Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn) Dist. Parbhani - 431511 (M.S.)




PRINCIPAL
Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn.) Dist. Parbhani 431511

Results:

Isolation and maintenance of Aeromonas hydrophila from various location of Marathwada region

A Total of eighteen (18) isolates of Aeromonas hydrophila were isolated from eight different

location of Marathwada districts. 04 isolates each were isolated from Nanded district, 02 isolates each were isolated from Parbhani, Hingoli, Latur, Beed, Osmanabad, Jalna and Aurangabad districts each (Table 1). These isolates were identified and maintained on their characteristics of growth on Nutrient agar slants.

Table 1: Isolate codes of Aeromonas hydrophila with their location of isolation

Sr. No	District	Natural culture system	Isolate code of <i>Aeromonas hydrophila</i>	Artificial culture system	Isolate code of <i>Aeromonas hydrophila</i>	Total
1	Nanded	Godavari (Vishnupuri)	Ah-01	Kandhar	Ah-03	04
		Asna (Devapur)	Ah-02	Petwadaj	Ah-04	
2	Hingoli	Kayadu (Balapur)	Ah-05	Bhategaon	Ah-06	02
3	Parbhani	Masoli (Gangakhed)	Ah-07	Yeldari	Ah-08	02
4	Aurangabad	Nagzhari (Tembapuri)	Ah-09	Paithan	Ah-10	02
5	Jalna	Galhati (Ambad)	Ah-11	Ghanewadi	Ah-12	02
6	Latur	Manjra (Latur)	Ah-13	Sakol, Udgir	Ah-14	02
7	Beed	Sindphana (patoda)	Ah-15	Mehakari, Ashti	Ah-16	02
8	Osmanabad	Sina (Paranda)	Ah-17	Khasapur	Ah-18	02
			09		09	18

Pathogenicity Studies of two Aeromonas (AH-04 and AH-07) isolates in fish ulcer disease using Cirrhinus mrigala

Two isolates Aeromonas (AH-04 and AH-07) were purified from the skin ulcers of the fish concerned. Since these bacteria were isolated from the fish's ulcer tissue, it was considered logical to study the clinical symptoms, histopathology and hematology of fish infected

with these pathogenic bacteria. The study of the pathogenicity and virulent characteristics of these bacteria were studied in detail.

Pathogenicity studies of the bacteria AH-04 and AH-07

The preliminary tests indicated that ulcers were associated with AH-04 and AH-07. It was thought proper to determine the amount of


Co-ordinator
IQAC

Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn) Dist. Parbhani - 431511 (M.S.)




PRINCIPAL 1727
Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn.) Dist. Parbhani

inoculum required to cause symptoms in healthy fishes. There are little records of these bacteria's LD50 values in any fish.

Therefore, experiments were conducted to establish the LD50 values of these bacteria, and a comparative study on the sensitivity of these fish species to these pathogenic bacteria was carried out.

Determination of the LD50 value

Bacteria isolates AH-04 and AH-07 cell suspensions were injected into healthy fish in serially diluted doses ranging from 1×10^2 to 1×10^7 cfu in 0.05 ml of inoculum. Materials

and Methods gives details of procedures followed. Fish mortality rates increased with all bacterial suspensions with increasingly concentrated. The bacterial isolates AH-04 and AH-07 could be purified from the dead fish lesion of liver and kidney. At the highest concentration (1×10^7 cfu), 85% of the fish inoculated with AH-04 died and AH-04 died at the lowest concentration (1×10^3 cfu). The LD50 values were estimated as 0.837×10^5 , and 0.878×10^5 for isolates AH-04 and AH-07, respectively (Table 2). The inoculums concentration 1×10^2 did not induced death or symptoms in fishes.

Table 2: Cumulative mortalities of *Cirrhinus mrigala* after 15 days of inoculation with serial 10 fold dilutions of AH- 04 and AH-07 (*Aeromonas hydrophila*).

Dose (c.f.u.)	number of fishes inoculated	Number of dead fishes	
		AH-04	AH-07
1×10^2	20	0	0
1×10^3	20	3	4
1×10^4	20	5	6
1×10^5	20	9	10
1×10^6	20	12	12
1×10^7	20	14	14
LD50		0.895×10^5	0.946×10^5

c.f.u 0.05 ml of inoculum

b Calculated number of bacteria required to kill 50% of injected fish. (Calculated from relationship between probits of percentage mortalities and the logs of the dilution series of bacterial suspension).

Comparative mortality rate of the two fish species

The mortality rate was 65.0% when inoculated with a mixed bacterial suspension of AH-04 and AH-07 in *Cirrhinus mrigala*. The pure bacterial suspensions, AH-04 induced 35 % and AH-07 induced 40 % mortality (Table 3).


Co-ordinator
IQAC
Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn) Dist. Parbhani - 431511 (M.S.)




PRINCIPAL
Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn.) Dist. Parbhani

Table 3: Percentage mortality and Nature of ulcer formation in Cirrhinus mrigala injected intramuscularly with saline suspensions of AH-04 and AH-07 in pure and mixed condition.

Bacterial strain	Number of fishes inoculated	Number of fishes dead	Nature of ulcer	Percentage mortality
Control	20	0	-	-
AH-04	20	9	moderate	45.0%
AH-07	20	11	moderate	55.0%
Mixed	20	14	severe	70.0%

Control set of fishes were intramuscularly injected with sterile saline suspension

Cumulative mortalities of fishes with respect to number of days after inoculation is shown in Table 4. The table shows that in all cases there is a steep rise in the number of mortalities after one after inoculation of pathogenic bacteria AH-04. The maximum number of death of fishes occurred during 4 to 7 days hours after

inoculation. After 7 days, the death rate fell steeply and no further death in the fishes were recorded. In Cirrhinus mrigala the death rate of fishes injected with AH-04 was slightly varied in fishes' species inoculated. The percentage mortality was higher in Cirrhinus mrigala.

Table 4 relation of time and Mortalities of the Cirrhinus mrigala after injection with bacterial suspension of AH-04

Days after inoculation	Number of fishes inoculated	AH-04		AH-07
		Number of fishes dead	% mortality	Number of fishes dead
1	20	0	0	0
2	20	2	10	3
3	20	7	35	7
4	20	9	45	10
5	20	11	55	12
6	20	12	60	13
7	20	13	65	16
8	20	14	70	16
9	20	15	75	16
10	20	15	75	16
11	20	15	75	16

**Ce-ordinator
IQAC**

Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn) Dist. Parbhani - 431511 (M.S.)



PRINCIPAL 1729
Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn.) Dist. Parbhani

12	20	15	75	16
----	----	----	----	----

Discussion:

In the present work, the serially diluted dose of different bacterial suspensions produced consistent trends of mortality. Isolation and identification of injected bacterial strains from external lesion, liver, and kidney of freshly dead fishes clearly indicate that the cause of death was associated with these bacterial strains. Various workers have observed changes in the liver and kidney of fishes infected by pathogenic *Aeromonas* strains (Pal and Pradhan, 1995). Prasad et al., (1995) could reisolate the pathogenic strain of *Aeromonas hydrophila* from liver, kidney and spleen of fish experimentally infected by the bacteria.

The LD₅₀ values of the two *Aeromonas* (AHS02 and AHS03) in fishes were found to be 1×10^5 c.f.u., for all four isolates. Descriptions of the degree of virulence by previous workers (Santos et al., 1996) suggested that strains of *Aeromonas* sp. showing LD₅₀ less than 1×10^6 could be classified as virulent, those with LD₅₀ 106 to 108 as weakly virulent and those with LD₅₀ more than 108 as nonvirulent. Accordingly AHS02 and AHS03 could be denoted as virulent. There was no significant difference between the virulence of these. The virulence was similar as suggested by its slightly higher LD₅₀ value.

Sahu et al., (1996) found that experimentally infected *Aeromonas hydrophila* could produce lesion in the catfish, *Clarias batrachus* on 30th day and in Rohu (*Labeo rohita*) on the 10th day. However, the infection was limited to ulcer in the skin and muscle only. Histopathological changes in the internal organs were not as remarkable as it was in the skin and muscle. Besides *Aeromonas* sp. and *Pseudomonas* sp.,

various other types of bacteria were also found to be associated with epizootic ulcerative syndrome. Jhingran and Das (1990) observed that *Micrococcus* sp. isolated from lesions and haematopoietic tissues of epizootic ulcerative syndrome affected fishes could induce ulcers within 72 hours both through inoculation and when kept in association with the bacteria. Kar et al., (1990) found *E. coli* and *Pseudomonas aeruginosa* to be associated with epizootic ulcerative syndrome affected fishes.

Ali and Tamuli (1991) isolated three types of bacteria from ulcers of four species of affected fishes and reinfection studies showed that *Aeromonas* sp. induced only mild infections, *Vibrio* sp. induced similar disease symptoms and *Micrococcus* sp. failed to induce any disease symptom.

Globally *Aeromonas* sp. is one of the most common bacteria associated with fish diseases. Although many strains are regarded as opportunistic pathogens, others are clearly primary pathogens in their own right (Trust, 1986). The fish diseases which involve *Aeromonas hydrophila* includes motile *Aeromonas* septicemia, red spot disease of European eel, *Anguilla anguilla* (Schaperclaus, 1934), red disease of Japanese eel, *A. japonica* (Hoshina, 1962) and red disease of carp, *Cirrhinus mrigala* (Egusa, 1978). Jo and Onishi (1980) isolated *A. hydrophila* from all diseased cultured ayu, *Plecoglossus altivelis* characterized by exophthalmus and subcutaneous ulceration. Rahim et al., (1985) isolated *A. hydrophila* from the wounds of five species of fishes in Bangladesh. Okpokwasili and Okpokwasili (1994) found that *Pseudomonas* spp. and *A. hydrophila* isolated from brown patch disease of tilapia were more

Co-ordinator
IQAC

Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn) Dist. Parbhani - 431511 (M.S.)



PRINCIPAL

Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn.) Dist. Parbhani

virulent to tilapia fingerlings when infected by a mixed culture than *A. hydrophila* or *Pseudomonas* spp. alone. Esteve et al., (1993) isolated *Aeromonas hydrophila* and *Aeromonas jandaei* from diseased European eel (*Anguilla anguilla*) from an eel farm in Spain which caused ulcerous disease by intraperitoneal injection (LD50 dose: 1×10^5 to 1×10^7 c. f. u. / fish) and also by bath exposure to 107 to 108 c.f.u./ ml in healthy eels.

Aeromonas liquifaciens, another pathogenic strain of *Aeromonas* sp. was isolated from scale protrusion disease in carp by Kusuda and Takahashi (1970), which affected fish farms of Japan. Reinfection studies in carp and gold fish showed that percentage mortality of carp was higher than that of gold fish. *Aeromonas punctata* has been regarded as the etiological agent of infectious dropsy in carps by Schaperclaus (1965) and he thought that primary infections resulting in disease may be induced by *Pseudomonas fluorescens*. Later he indicated that 3 different microorganism, *A. punctata*, *P. fluorescens* and a virus may be involved in this disease which manifest itself in several forms (Schaperclaus, 1969).

This tends to indicate that *Aeromonas* sp. are highly opportunistic pathogens which invade the fish once the skin barrier is breached. This however does not eliminate the fact that these bacteria are primary pathogens. In order to arrive at a conclusion on this aspect, a detailed study on the role of these bacteria in causing epizootic ulcerative syndrome is necessary.

Reference

Ali, A. and K. Tamuli. 1991. Isolation of an etiological agent of epizootic ulcerative syndrome. *Fish. Chimes*. 11: 43-46.

Barde, R. D. (2021). Bacterial Diversity Associated With *Labeo Rohita* Isolated From Cultured And Natural Water Bodies.

European Journal of Molecular & Clinical Medicine, 7(11), 2020.

Barde, R. D., Apastambh, A. R., & Baig, M. M. V. (2021). Association of Oomycetes *Aphanomyces invadans* with Freshwater fishes. *Annals of the Romanian Society for Cell Biology*, 25(6), 14118-14125.

Barde, R. D., Deshpande, M., Nagthane, N., Darak, O., & Baig, M. M. V. (2020). A Review of Saprolegnia Infection in Freshwater Fishes and Control of the Saprolegniosis. *Sustainable Humanosphere*, 16(1), 702-711.

Bergey's manual of Systematic Bacteriology 1986. Vol.II (Eds.A. Peter, A. Sneath and M. Elizabeth sparbe). Williams and Wikins, Baltimore. pp. 156-170.

Cheesbrough, M. 1989. Medical laboratory manual for tropical countries, Microbiology, Vol.II, Ed. Tropical health technology, Butterworth and Co.Ltd.,Cambridge, pp. 216-252.

Darak, O. S., & Barde R.D., (2014). Association of bacterial pathogens with *Labeo rohita* in Marathwada region of Maharashtra. *Int. J. Adv. Biotechnol. Res.* 15, 777-783.

Darak, O. S., Baisthakur, P. O., & Barde, R. D. (2021). Antibiotic Resistance in *Micrococcus* SP Isolated from Freshwater Fish Farms *Catla Catla*. *EFFLATOUNIA-Multidisciplinary Journal*, 5(2).

Darak, O., & Barde, R. D. (2015). *Pseudomonas fluorescens* associated with bacterial disease in *Catla catla* in Marathwada region of Maharashtra. *International Journal of Advanced Biotechnology and Research*, 6(2), 189-195.

Egusa, S. 1978. Infectious diseases of fishes Koseisha-Koseikaku, Tokyo, Japan. pp.

Co-ordinator
IQAC

Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn) Dist. Parbhani - 431511 (M.S.)



PRINCIPAL 1731
Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn.) Dist. Parbhani

- Esteve, C., E. G. Biosca and C. Amaro. 1993. Virulence of *Aeromonas hydrophila* and some other bacteria isolated from European eels, *Anguilla anguilla* reared in fresh water. *Dis. Aquat. Org.* 16: 15-20.
- Hoshina, R. 1962. Studies on red disease of Japanese eel. *J. Tokyo Univ. Fish.* 6: 1-104.
- Jhingran, A. G. and M. K. Das, 1990. Epizootic ulcerative syndrome in fishes. Bull. No. 65. Central Inland Capture Fisheries Research Institute, Barrackpore, India.
- Jo, Y. and K. Onishi. 1980. *Aeromonas hydrophila* isolated from cultured Ayu (*Plecoglossus altivelis*). *Fish Pathol.* 15: 85-90.
- Kar, D., R. G. Michael, S. Kamad and S. Changkija. 1990. Studies on fish epidemics from Assam. *J. Inland Fish Assoc.* 20: 73-75.
- Kusuda, R. and Takahashi, Y., 1970. Studies on the Scale Protrusion Disease of Carp Fishes-I Characteristics of *Aeromonas liquefaciens* Isolated from Diseased Fishes. *Fish Pathology*, 4(2), pp.87-97.
- Okpokwasili, G. C. and N. P. Okpokwasili. 1994. Virulence and drug resistant patterns of some bacteria associated with "ubrown patch" disease of tilapia. *J. Aqua. Trop.* 9: 223-233.
- Pal, J. and K. Pradhan. 1995. Histopathology of liver, kidney and spleen of experimentally infected fish, *Clarias batrachus*. Fourth Asian Fisheries Forum, Abstracts, Oct. 16-20, 1995, Beijing, China. pp. 46.
- Prasad, Y., T. A. Qureshi, S. A. Mastan and R. Chouhan. 1995. Studies on the pathogenicity of *Aeromonas hydrophila* isolated from EUS affected fishes. Fourth Asian Fisheries Forum, Abstracts. October 16-20, 1995, Beijing, China. pp. 38.
- Rahim, Z., K. M. S. Aziz, M. I. Haque and H. Saeed. 1985. Isolation of *Aeromonas hydrophila* from the wounds of five species of brackishwater fish of Bangladesh. *Bangladesh J. Zoo.* 13: 37-42.
- Sahu, B. B., Gayatri Murjani and S.C. Mukherjee. 1996. *Aeromonas* infection in freshwater fish and catfish. National workshop on fish and prawn disease epizootics and quarantine adoption on India. Oct. 9, 1996. Central Inland Capture Fisheries Research Institute, Barrackpore, India. pp 87-93.
- Santos, Y. Bandin, I. and Toranzo, A. E. 1996. Immunological analysis of extracellular products and cell surface components of motile *Aeromonas* isolated from fish. *J. Appl. Bacteriol.* 81: 585-593.
- Schaperclaus, W. 1934. Untersuchungen uber dielseuchen in deutschen kinnerund kusten wasseren 1930-1933. *Zeit. Fisch.* 32: 191-217.
- Schaperclaus, W. 1969. Virus infektionen befischen. In *Handbuch der virus infektionen bei Tieren*. (H. Rohrer, ed.) pp. 1067-1141. VEB Gustav. Fischer, Jena.
- Trust, T. J. 1986. Pathogenesis of infectious diseases of fish. *Ann. Rev. of Microbiol.*


Co-ordinator
IQAC
Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn) Dist. Parbhani - 431511 (M.S.)




PRINCIPAL
Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn.) Dist. Parbhani