



Study of macromolecular contents of different tissues of *Channa punctatus* under stress condition.

Thakur PS¹ and Chavan PN^{2*}

¹Department of Zoology, Shri Shivaji College, Akola. (MS), India

²Department of Zoology, S.G.B. Mahavidyalaya, Purna, (MS), India

*Corresponding Author email: pnychavan.2009@rediffmail.com

Manuscript details:

Received: 13.02.2020

Accepted: 09.03.2020

Published: 10.04.2020

Cite this article as:

Thakur PS and Chavan PN (2020) Study of macromolecular contents of different tissues of *Channa punctatus* under stress condition, *Int. J. of Life Sciences*, Volume 8(1): 168-170.

Copyright: © Author, This is an open access article under the terms of the Creative Commons Attribution-Non-Commercial - No Derives License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Available online on <http://www.ijlsci.in>
ISSN: 2320-964X (Online)
ISSN: 2320-7817 (Print)


ABSTRACT

Channa has a wide natural distribution extending from Iran in the west, to China in the east, and parts of Siberia in the Far East. They are one of the most common staple food fish in Thailand, Cambodia, Vietnam and other South East Asian countries where they are extensively cultured. Apart from their importance as a food fish, snakeheads are also consumed as a therapeutic for wound healing as well as reducing post-operative pain and discomfort, and collected for the international aquarium pet trade. The stress response is complex and involves a series of behavioral and physiological adaptive response which enables the fish to overcome the effects of stress. These responses involve all levels of animal organization and is collectively called integrated stress response. Present study has been undertaken to examine the changes in fish, *Channa punctatus* after chronic exposure to such levels of stress due to starvation in laboratory condition, which are normally present in water bodies around Akola.

Keywords : *Channa punctatus* , stress response, chronic exposure.

INTRODUCTION

Channa is a genus of the Channidae family of snakehead fish. It has a great demand in market because of its high nutritional value. Study of food and feeding habits of fishes have manifold importance in fishery biology. Seasonal change in temperature not only influence food consumption and rate of digestion but also quality and quantity of available food organisms. Studies on the food and feeding habits of different fishes have been made by different workers (Bulow'1970, Collvin *et al.* 1985). Food stress and space stress has been recognized to play a vital role in almost all aspects of living systems either directly or indirectly (Dhawan and Kaur, 1997; Sastry and Shukla *et al.*, 1994). However, very few reports are available on the extent of changes in the nutritive value of fish after prolonged exposures to stress. The stress has been defined as response of the cell or organism to any demand placed on it which causes an extension of the physiological state beyond the normal resting state. The stress response is complex and involves a series of



PRINCIPAL
168

behavioural and physiological adaptive response which enables the fish to overcome the effects of stress. These response involve all levels of animal organization and is collectively called integrated stress response. Under natural conditions numerous fish species endure long periods of starvation associated mainly with seasonal changes in food.

MATERIAL AND METHOD

The samples for the present study were collected from the local market, Mature *Channa punctatus* weighing from 150-200 gm and mean length 27 ± 3 cm were procured and acclimated at $24.5^{\circ}\text{C} - 35.5^{\circ}\text{C}$ for 15 days. (Shrama and Davis, 1980) The natural photoperiod was maintained during the period. They were acclimatized to laboratory conditions for 2 weeks. Thirty individuals were placed in aquarium for 60 days, from end of November 2011 to end of January 2012. The aquarium having 40 liters of Tap water which was dechlorinated (pH 7.0- 7.9) at least for experimentation. They were fed alternate days, regularly during acclimatization period. After

acclimatization for two weeks, healthy specimens of *Channa punctatus* were divided into control and fasting groups. Each group contained 10 individuals. Control was fed on chopped earthworms. However, the food was withheld from the fasting group. Liver, Muscle, Gill were immediately dissected out for estimation of DNA and RNA (Sornarej 1995). The nucleic acid content was determined by the method of biochemical analysis (Volkin E. and Cohn W.E. 1967).

RESULTS AND DISCUSSION

It is evident from the starvation studies that the fasting decreases significantly the activity of fish. Starvation resulted in changes in body weight and chemical composition (Sunnep *et al* 2011). Muscle rich in proteins, forms mechanical tissue intended for mobility and do not participate in metabolism. Liver being the centre for various metabolisms is also rich in proteins. And DNA contents is responsible for the formation of protein molecules. Decrease in DNA and RNA content of muscles and liver in stress, decrease in DNA and RNA content in gills.




Figure 1:- Variation in the DNA contents (mg/g) of the liver , Gills and Muscle of *Channa punctatus*




Figure 2:- Variation in the RNA contents (mg/g) of the liver, Gill and Muscle of *Channa punctatus*.

The values of result were represented by means of standard deviations. It is evident from the starvation studies that the fasting decreases significantly the activity of fish. Starvation is experienced in most species of fish during certain periods of every year largely due to environmental conditions (Mali and Chavan 2011). The concentration of DNA and RNA depicted that affect the concentration of glycogen and protein from the different tissues of *Channa punctatus*.

SUMMARY AND CONCLUSION

Around Akola region only some small water bodies are their in which *Channa punctatus* cultured naturally. But during summer, environmental temperature go rise up to 48°C, that's why water bodies become start to convert into dry lands, naturally aquatic animals gathered in a small place available for to sustained there life. Fish like *Channa punctatus* getting starve and under stress due to the small water available for sustained there life (Sunnap *et al.*, 2011). Gills, muscles and liver are the important organ of fish *Channa punctatus* for to survive, and all the activity of these organs depends upon the protein available for to produce energy and DNA is responsible for to produce the protein molecules. DNA bound inside the cell with the help of nuclear membrane. During the stress, some enzymes are responsible for the shading of cells ultimately responsible for the loss in genetic material.

Conflict of Interest

The author declares that there is no conflict of interest.

REFERENCES

- Bulow FJ (1970) RNA-DNA ratios as indicators of recent growth rates of a fish. *J. Fish. Res. Board Can.*, 27: 2343-2349.
- Collvin I (1985) Effects of copper on growth and starvation in perch, *Perca fluviatilis*. *J. Fish. Biol.*, 27(6): 757-764.
- Dhawan A and Kaur K (1997) Effect of zinc on maturation and breeding potential of *Channa punctatus* and *Cirrhina mrigala*. *Int. J. Environ. Stud.*, 53: 265-274.
- Mali RP and Chavan PN (2011) Starvation and its impact on biochemical aspects of fresh water fish *Oreochromis mossambicus*.
- Sastry KV and Shukla V (1994) Acute and chronic toxic effects of cadmium on some haematological, biochemical and enzymological parameters in the fresh water teleost *Channa punctatus*. *Acta. Hydrochim. Hydrobiol.*, 22(4): 171-176.

Sharma DC and Davis PS (1980) Effect of methyl mercury on protein synthesis in liver of European carp, *Cyprinus carpio*. *Ind. J. Exp. Biol.*, 18(9): 1054-1055.

Sornarej R, Thanalashmi S and Bhaskaran P (1995) Influence of heavy metals on biochemical responses of the freshwater air breathing fish *Channa punctatus* (Bl). *Ecotoxicol. Environ. Monit.*, 5(1): 19-27.

Sunnap N, Nanware S and Chavan P (2011) Effect of starvation on the biochemical composition of freshwater fish *Channa punctatus*. *Recent Research in Science and Technology* 3 (9) 17-19.

Volkin E and Cohn WE (1967) Estimation of nucleic acids, In method of biochemical analysis, edited by D. Glick, Vol 1 pp 287-365.

© 2020 | Published by IJLSCI