



SYNTHESIS OF POLYHYDROQUINOLINES USING PHENYL PHOSPHONIC ACID AS CATALYST

Sanjeev M. Reddy¹ and Jitendra S. Pulle^{2*}¹Department of Chemistry, G.M.V. Kotgryal, Dist. Nanded (M.S.) India.²Department of Chemistry, S.G.B. College, Purna (Jn.), Dist. Parbhani (M.S.), India.

*Corresponding Author: Dr. Jitendra S. Pulle

Department of Chemistry, S.G.B. College, Purna (Jn.), Dist. Parbhani (M.S.), India.

Article Received on 26/01/2020

Article Revised on 16/02/2020

Article Accepted on 06/03/2020

ABSTRACT

A facile and efficient method has been developed for the one pot synthesis of polyhydroquinolines *via* four component reaction of ethyl acetoacetate, aldehyde, dimedone and ammonium acetate in the presence of catalytic amount of phenyl phosphonic acid in ethanol through Hantzsch reaction. The method described herein is simple, environmentally benign and provides good yields of the products.

KEYWORDS: Polyhydroquinolines, phenyl phosphonic acid, ethyl acetoacetate, aldehyde, dimedone.

INTRODUCTION

1,4-Dihydropyridines (DHPs) are important class of nitrogen containing compounds which received much attention due to their wide range of pharmaceutical and biological properties.^[1] They are found in biologically active products such as vasodilator, bronchodilator, antitumor, hepatoprotective, geroprotective, anti-atherosclerotic and antidiabetic agents.^[2] 1,4-DHP compounds have their important role in medicinal chemistry as calcium channel blockers as exemplified by therapeutic agents such as nifedine, nitrendipine, amlodipine, felodipine and nicardipine which are the important drugs used in cardiovascular disease.^[3] Recent studies have revealed that 1,4-DHPs exhibit several medicinal applications which include neuroprotectant^[4] and plate anti aggregatory activity.^[5] Also they have been reported for their applications in treatment of Alzheimer disease due to their cerebral antischematic activity and chemosensitizer in tumor-therapy.^[6] These properties and applications clearly show the potential of 1,4-DHPs as valuable drug candidates.

In 1882, Arthur Hantzsch reported first synthesis of substituted 1,4-dihydropyridine by one pot condensation of ethyl acetoacetate, aromatic aldehyde, dimedone and ammonia. The reaction was conducted in acetic acid or at reflux in ethanol for long periods resulting in low to moderate yield of products.

Owing to the wide range of biological and medicinal applications, the synthesis of such compounds has received much attention in recent years. Many synthetic


methods have been reported for the preparation of polyhydroquinoline derivatives. Some of the techniques, catalysts and reagents such as Microwave^[7], Solid phase organic synthesis technique^[8], Molecular iodine^[9], TMSCl^[10], HClO₄-SiO₂^[11], Cerium (IV) ammonium nitrate^[12], L-Proline^[13], Without catalyst^[14], Ionic liquids^[15], Polymers^[16], Alumina sulphuric acid^[17], ZnCl₂^[18], Iron (III) trifluoroacetate^[19], Heteropoly acids^[20], *p*-Toluene sulphonic acid^[21], Organo catalyst^[22], 3-Nitrophenyl boronic acid^[23], Copper perchlorate hexahydrate^[24], Metal triflates^[25], Nafion-H^[26], Aluminum phosphate [AlPO₄(H)]^[27], Palladium (0) nanoparticles^[28], ZnO-beta zeolite^[29], Hg- II Complex^[30], Cu -II Complex^[31], MNPs -TBSA^[32], Nicotinic acid^[33], have been used for the synthesis of polyhydroquinolines. Each of these above methods have their own merits while some of the methods suffer from long reaction times, low yields, harsh reaction conditions, tedious work ups, etc. For these reasons, efforts have been made to replace such conventional catalysts by other catalysts. The research activities and industrial sections now a days require development of catalysts^[34] and the eco-friendly catalytic synthesis which offer strong potential for fast application in industries.^[35]

We herein report phenyl phosphonic acid as an efficient Bronsted acid catalyst for the synthesis of polyhydroquinolines V in the condensation reaction of β -keto ester III, ammonium acetate IV, dimedone II and various aromatic aldehydes I in ethanol at room temperature (Scheme 1).

Co-ordinator
IQACShri Guru Budhivamsi Mahavidyalaya
Purna, Dist. Parbhani - 431512 (M.S.)
www.ejpmr.com

PRINCIPAL

Shri Guru Budhivamsi Mahavidyalaya
Purna, Dist. Parbhani - 431512 (M.S.)




Scheme 1.

Experimental procedure

A mixture of dimesone (1 mmol), aromatic aldehyde (1 mmol), ethyl acetoacetate (1 mmol), ammonium acetate (1 mmol) and phenyl phosphonic acid (30 mg) was stirred in ethanol (3 mL) for 2-3 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was poured into ice cold water and stirred for about 10-15 minutes. The product was then filtered, dried and recrystallized from ethanol to get pure polyhydroquinoline derivatives.


Spectral analysis

The products obtained were characterized by their IR, ¹H NMR and Mass spectra. The analysis of some of the representative compounds is given below.



Ethyl-1, 4, 7, 8-tetrahydro-2, 7,7-trimethyl-4(phenyl)-5-(6H)-oxoquinolin-3-carboxylate

IR (cm⁻¹) : 3286(-NH-), 1697(C=O), 1608 (C=O)
¹H NMR (δ) : 0.98(s,3H,CH₃), 1.05(s,3H,CH₃), 1.15(t,3H,CH₃), 2.13-2.36(m,4H,2x CH₂), 2.4(s,3H,CH₃), 7.1-7.4(m, 5H,Ar-H)
Mass : 328 (M⁺)



Ethyl-1, 4, 7, 8-tetrahydro-2, 7,7-trimethyl-4(4-chlorophenyl)-5-(6H)-oxoquinolin-3-carboxylate

5-(6H)-oxoquinolin-3-carboxylate

IR (cm⁻¹) : 3276(-NH-), 1705(C=O), 1605 (C=O)
¹H NMR (δ) : 0.92(s,3H,CH₃), 1.07(s,3H,CH₃), 1.15(t,3H,CH₃), 2.13-2.35(m,4H,2xCH₂), 2.38(s,3H,CH₃), 4.05(q,2H,CH₂), 5.02(s,1H,CH), 5.5(s,1H,NH), 7.12-7.28(m,4H,Ar-H)
Mass : 363 (M⁺)

Table 1: Synthesis of polyhydroquinoline derivatives using phenyl phosphonic acid.

Entry	Aldehyde	Product	Time (hrs)	Yield (%)	M.P.(°C) [Observed]	M.P.(°C) [Reported]
1			3	92	203-205	202-204 [13]
2			2.5	92	230-232	232-234 [13]
3			2	89	207-208	208-210 [15a]









Co-ordinator
IQAC












Shri Guru Buddhiswami Mahavidyalaya
Purna (Jr) Dist. Parbhani - 431511 (M.S.)
www.ejpmr.com



PRINCIPAL

Shri Guru Buddhiswami Mahavidyalaya
Purna (Jr.) Dist. Parbhani

4			3	90	242-243	241-243 [25b]
5			3	91	260-262	-
6			3	90	288-290	-
7			3	93	108-110	-

Entry	Aldehyde	Product	Time (hrs)	Yield (%)	M.P. (°C) [Observed]	M.P. (°C) [Reported]
8			2.5	92	240-242	241-243 [13]
9			3	90	182-184	-
10			2.5	92	252-253	253-255 [25b]
11			2.5	91	265-266	-
12			2	89	200-202	-
13			2	90	232-234	233-234 [13]

Co-ordinator
IQAC

RESULTS AND DISCUSSION

The studies of facile four-component Hantzsch condensation reactions in presence of phenyl phosphonic acid catalyst at room temperature were explored. The results are summarized in Table 1. The condensation of aldehydes, dimedone, ethyl acetoacetate and ammonium acetate was found to be effective within 2-3 hours in ethanol to produce the corresponding pharmacologically useful polyhydroquinoline derivatives in good yields. Various aromatic aldehydes having electron-donating and electro-withdrawing substituent underwent the smooth reaction. It is noteworthy that the structural variation of the aldehydes and substituent on the aromatic ring did not show any effect on the conversion, as the products were obtained in good yields in short reaction times at mild reaction conditions. Thus phenyl phosphonic acid, Bronsted acid catalyst was found to be an efficient catalyst for the Hantzsch polyhydroquinoline derivatives.

CONCLUSION

Polyhydroquinoline derivatives via Hantzsch condensation were obtained in good yields using phenyl phosphonic acid as a Bronsted acid promoter. The method was found to be suitable with various functional groups under mild conditions in one-pot condensation.

REFERENCES

- Kato M, Nakajima M, Shimada N, Yamazaki H, Yokoi T. Inhibition of human cytochrome P450 enzymes by 1,4-dihydropyridine calcium antagonists: prediction of in vivo drug-drug interactions. *Eur. J. Clin. Pharmacol.* 2000; 55: 843-2.
- Shan R, Valazquez C, Knaus E. Syntheses, Calcium Channel Agonist-Antagonist Modulation Activities, and Nitric Oxide Release Studies of Nitrooxalkyl 1,4-Dihydro-2,6-dimethyl-3-nitro-4-(2,1,3-benzoxadiazol-4-yl) pyridine-5-carboxylate Racemates, Enantiomers, and Diastereomers. *J. Med. Chem.* 2004; 47(1): 254-1.
- (a) Ortiz ME, Nunez - Vergara LJ, Sequella JA. Relative Reactivity of Dihydropyridine Derivatives to Electrogenated Superoxide Ion in DMSO Solutions: A Voltammetric Approach. *Pharm. Research*, 2003; 20: 292-6; (b) Budriesi R, Bisi A, Ioan P, Rampa A, Gobbi S, Belluti F, Piazzi L, Valenti P, Chiarini A. 1,4-Dihydropyridine derivatives as calcium channel modulators: the role of 3-methoxy-flavone moiety. *Bio organic and Med. Chem.* 2005; 13(10): 3423-0; (c) Miri R, Javidnia K, Sarkaradeh H, Hemmetenejad B. Synthesis, study of 3D structures, and pharmacological activities of lipophilic nitroimidazolyl-1,4-dihydropyridines as calcium channel antagonist. *Bio organic and Med. Chem.* 2006; 14(14): 4842-9.
- Bretzel RG, Bollen CC, Maeser E, Federlin KF. Trombopline platelet aggregation inhibitor antithrombotic. *Drugs Fut*, 1992; 17: 465-9.
- Boer R, Gekeler V. Chemosensitizers in tumor therapy: new compounds promise better efficacy. *Drugs Fut*, 1995; 20(5): 499-9.
- (a) Hantzsch A. Ueber die Synthese pyridinartiger Verbindungen aus Acetessigäther und Aldehydammoniak. *Justus Liebig's Annalen der Chemie.* 1882; 215(1): 1-2; (b) Hantzsch A. Synthese von Thiazolen und Oxazolen. *Ber. Dtsch. Chem. Ges.*, 1888; 21(1): 942-6; (c) Hantzsch A. Ueber das sogenannte Cyanacetone. *Ber. Dtsch. Chem. Ges.* 1890; 23(1): 1472-4; (d) Wiley R H, England D C, Behr LC. *Inorganic Reactions*. Wiley: Toronto, 1951; 16: 367.
- Ohberg L, Westman J. An Efficient and Fast Procedure for the Hantzsch Dihydropyridine Synthesis under Microwave Conditions. *Synlett*, 2001; 8: 1296-8.
- Gordeev MF, Patel DV, Gordon PM. Approaches to Combinatorial Synthesis of Heterocycles: A Solid-Phase Synthesis of 1,4-Dihydropyridines. *J. Org. Chem.* 1996; 61(3): 924-8.
- (a) Ko S, Sastry MNV, Lin C, Yao C F. Molecular iodine-catalyzed one-pot synthesis of 4-substituted-1,4-dihydropyridine derivatives via Hantzsch reaction. *Tetrahedron Lett.* 2005; 46(34): 5771-4; (b) Zelfigol MA, Salehi P, Zad AK, Shayegh M. Iodine-catalyzed synthesis of novel Hantzsch N-hydroxyethyl 1,4-dihydropyridines under mild conditions. *J. Mol. Catalysis A. Chem.* 2007; 261(1): 88-2.
- Sabitha G, Reddy G S, Reddy CS, Yadav JS. A novel TMSI-mediated synthesis of Hantzsch 1,4-dihydropyridines at ambient temperature. *Tetrahedron Lett.* 2003; 44(21): 4129-1.
- (a) Mahaswara M, Siddaiah V, Damu G LV, Rao CV. An efficient one-pot synthesis of polyhydroquinoline derivatives via Hantzsch condensation using a heterogeneous catalyst under solvent-free conditions. *Arkivoc.* 2006; ii: 201-6; (b) Mahaswara M, Siddaiah V, Rao YK, Tzeng YM, Sridhar C. A simple and efficient one-pot synthesis of 1,4-dihydropyridines using heterogeneous catalyst under solvent-free conditions. *J. Mol. Catalysis A. Chem.* 2006; 260(1-2): 179-0.
- (a) Ko S, Yao CF. Cerium Ammonium Nitrate (CAN) catalyzes the one-pot synthesis of polyhydroquinoline via the Hantzsch reaction. *Tetrahedron*, 2006; 62(31): 7293-9; (b) Reddy C S, Raghu M. Cerium (IV) ammonium nitrate catalyzed facile and efficient synthesis of polyhydroquinoline derivatives through Hantzsch multicomponent condensation. *Chin. Chem. Lett.* 2008; 19(7): 775-9.

Co-ordinator
IQAC

Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn) Dist. Purbhoni-431511 (N.S.)



PRINCIPAL
Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn) Dist. Purbhoni

13. Karade NN, Budhewar VH, Shinde S V, Jadhav W N. L-Proline as an Efficient Organo-Catalyst for the Synthesis of Polyhydroquinoline Via Multicomponent Hantzsch Reaction. *Lett. Org. Chem.*, 2007; 4(1): 16-9.
14. Arumugam P, Perumal PT. Hantzsch synthesis of polyhydroquinolines - A simple, efficient and neat protocol. *Indian J. Chem. Sec. B*, 2008; 47: 1084-0.
15. (a) Ji SJ, Jiang ZQ, Lu J, Loa TP. Facile Ionic Liquids-Promoted One-Pot Synthesis of Polyhydroquinoline Derivatives under Solvent Free Conditions. *Synlett*, 2004; 5: 831-5; (b) Sridhar R, Perumal PT. A new protocol to synthesize 1,4-dihydropyridines by using 3,4,5-trifluorobenzeneboronic acid as a catalyst in ionic liquid: synthesis of novel 4-(3-carboxyl-1H-pyrazol-4-yl)-1,4-dihydropyridines. *Tetrahedron*, 2005; 61(9): 2465-0.
16. Breitenbucher JG, Figliozzi G. Solid-phase synthesis of 4-aryl-1,4-dihydropyridines via the Hantzsch three component condensation. *Tetrahedron Lett.*, 2000; 41(22): 4311-5.
17. Arslan M, Faydali C, Zengin M, Kucukislamoglu M, Demirhan, H. An efficient one pot synthesis of 1,4-dihydropyridines using alumina sulfuric acid (ASA) catalyst. *Turk. J. Chem.*, 2009; 33: 769-4.
18. Sanjeeva Reddy Ch, Raghu M. Facile $ZrCl_4$ promoted four-component coupling one-pot synthesis of polyhydroquinoline derivatives through unsymmetric Hantzsch reaction. *Indian J. Chem. Sec. B*, 2008; 47: 1578-2.
19. Adibi H, Samimi HA, Beygzadeh M. Iron(III) trifluoroacetate and trifluoromethanesulfonate: Recyclable Lewis acid catalysts for one-pot synthesis of 3,4-dihydropyrimidinones or their sulfur analogues and 1,4-dihydropyridines via solvent-free Biginelli and Hantzsch condensation protocols. *Catal. Commun.*, 2007; 8(12): 2119-4.
20. Heravi MM, Bakhtiri K, Javadi NM, Bamoharram FF, Saeedi M, Oskooi HA. $K_2[PW_{11}CoO_{40}]$ -catalyzed one-pot synthesis of polyhydroquinoline derivatives via the Hantzsch three component condensation. *J. Mol. Catalysis A. Chem.*, 2007; 264(1-2): 50-2.
21. Cherkupally SR, Mekala R. P-TSA Catalyzed Facile and Efficient Synthesis of Polyhydroquinoline Derivatives through Hantzsch Multi-Component Condensation. *Chem. Pharm. Bull.*, 2008; 56(7): 1002-4.
22. Kumar A, Maura RA. Synthesis of polyhydroquinoline derivatives through unsymmetric Hantzsch reaction using organocatalysts. *Tetrahedron*, 2007; 63(9): 1946-2.
23. Adude RN, Tigote RM, Goswami SV, Bhusare SR. 3-Nitrophenylboronic acid-catalyzed efficient one-pot synthesis of 1,4-dihydropyridines and polyhydroquinolines. *Int. J. Ind. Chem.*, 2012; 3(6): 1-3.
24. Puri S, Kaur B, Parmar A, Kumar H. Copper Perchlorate Hexahydrate: An Efficient Catalyst for the Green Synthesis of Polyhydroquinolines under Ultrasonication. *ISRN Organic Chemistry*, 2011; 1-4.
25. (a) Donelson JL, Gibbs RA, De SK. An efficient one-pot synthesis of polyhydroquinoline derivatives through the Hantzsch four component condensation. *J. Mol. Cat. A. Chem.*, 2006; 256(1-2): 309-1; (b) Wang LM, Sheng J, Zhang L, Han JW, Fan ZY, Tian H, Qian CY. Facile $Yb(OTf)_3$ promoted one-pot synthesis of polyhydroquinoline derivatives through Hantzsch reaction. *Tetrahedron*, 2005; 61(6): 1539-3.
26. Kidwai M, Chauhan R, Bhatnagar D, Singh AK, Mishra B, Dey S. Nafion-H catalyzed synthesis of polyhydroquinolines via the Hantzsch multi-component reaction. *Monatshefte für Chemie*, 2012; 143: 1675-0.
27. Purandhar K, Jyothi V, Pratap Reddy P, Adharvana Chari M, Mulkanti K. Aluminum phosphate $[AlPO_4(H)]$: A mild and efficient recyclable catalyst for one-pot synthesis of polyhydroquinoline via the Hantzsch reaction under solvent-free conditions. *Journal of Heterocyclic Chemistry*, 2012; 49(1): 232-6.
28. Saha M, Pal AK. Palladium(0) nanoparticles: an efficient catalyst for the one-pot synthesis of polyhydroquinolines. *Tetrahedron Letters*, 2011; 52(38): 4872-7.
29. Katkar S S, Mohite PH, Gadekar LS, Arbad BR, Lande MK. ZnO-beta zeolite: as an effective and reusable heterogeneous catalyst for the one-pot synthesis of polyhydroquinolines. *Green Chemistry Letters and Reviews*, 2010; 3(4): 287-2.
30. Ghasemi H D, Allameh S, Beyramabadi SA, Behmadi H, Morsali A. New Hg (II) complex: As a novel, green and efficient reusable catalyst for synthesis of polyhydroquinolines via Hantzsch condensation. *Asian J. Biol. Lif. Sci.*, 2018; 7(1): 17-0.
31. Raouf H, Allameh S, Beiramabadi SA. An Efficient and Facile Synthesis of Polyhydroquinolines through Hantzsch Reaction Catalyzed by a Novel and Reusable Cu (II) Complex. *J. Biochem Tech.*, 2018; 9(1): 61-4.
32. Bodaghifard MA. Bis Sulfamic Acid Functionalized Magnetic Nanoparticles as a Retrievable Nanocatalyst for the Green Synthesis of Polyhydroquinolines and Tetrahydrobenzopyrans. *J. Nanostruct.*, 2019; 9(1): 29-0.
33. Davarpanah J, Ghahremani M, Najafi O. Synthesis of 1,4-dihydropyridine and polyhydroquinoline derivatives via Hantzsch reaction using nicotinic acid as a green and reusable catalyst. *J. Mol. Struct.*, 2019; 1177: 525-5.
34. (a) Shanmugam S, Viswnathan B, Varadarajan, TK. Esterification by solid acid catalysts—a comparison. *J. Mol. Cat. A. Chemical*, 2004; 223(1-2): 143-7; (b) Amaranth CA, Palaniappan S, Saravanan C. Synthesis of benzyl acetate using polyaniline salts

Co-ordinator

IQAC

TQAC



- as catalysts - Part II. Poly. Adv. Tech, 2004; 15: 118-1.
35. Palaniappan S, John A. A novel polyaniline-fluoroboric acid-dodecylhydrogensulfate salt: versatile reusable polymer based solid acid catalyst for organic transformations. J. Mol. Cat. A: Chemical, 2005; 233(1-2): 9-5.



**Co-ordinator
IQAC**

Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn) Dist. Parbhani - 431511 (M.S.)



PRINCIPAL

Shri Guru Buddhiswami Mahavidyalaya
Purna (Jn.) Dist. Parbhani